RESIDENT VULTURES OF OMAN

EGYPTIAN VULTURE

(Rakhma) رخمة
Neophron percnopterus

Estimated global population: 18,600-54,000, declining
Number of subspecies: 3
IUCN* status: Endangered
Oman status: Breeding resident, few winter visitors



BREEDING BIOLOGY

- Monogamous.
- Breeds between February and May.
- Nests on cliffs, in shaded crevasses, ledges or small caves.
- Pairs maintained across years.
- Two-young clutches, many pairs fledge two young.
- Incubation averages 42 days.
- Young fledge at 70-85 days.

FEEDING BEHAVIOUR

- Feeds on carrion, waste from dumps and slaughterhouses, bird eggs, animal faeces and small living prey.
- Parents feed nestlings via regurgitation.
- Young follow parents to learn social skills and locate food.

MOVEMENT

- Wide ranging.
- Partial migrants.
- Breeds in Eurasia and Africa, migrants winter in Africa and southern Asia.

SOCIAL BEHAVIOUR

- Gregarious (sociable).
- Territorial breeding.
- Communal roosting.
- Group foraging (food finding).
- Human commensal (benefits from humans).



LAPPET-FACED VULTURE

(Nesr Al Athen) النسر الآذن
Torgos tracheliotos

Estimated global population: 9,200, declining
Number of subspecies: 2
IUCN* status: Endangered
Oman status: Breeding resident



BREEDING BIOLOGY

- Monogamous.
- Breeds between November and May.
- Nests on top of flat-topped trees.
- Parents share incubation, chick-rearing and shading.
- Single-egg clutch is the norm.
- Incubation averages 55-56 days.
- Young fledge at 105-120 days with parental care exceeding 3 additional months.

FEEDING BEHAVIOUR

- Feeds almost exclusively on carrion, can also take termites and locusts.
- Dominates other species at carcasses.
- Consumes skin, tendons and ligaments.
- Can also prey on eggs and nestlings.
- Routinely visits, bathes and drinks at watering holes.

MOVEMENT

- Wide ranging.
- Seasonal movement for some individuals but adults mostly sedentary.

SOCIAL BEHAVIOUR

- Relatively solitary.
- Territorial breeding.
- Communal roosting.
- Group foraging (food finding).
- Pairs often fly together searching for food, and perch together.



Did you know that Old World and New World Vultures have different taxonomies?

LEARN MORE

Find out more about raptors and their status in Oman here!

MAIN THREATS IN OMAN

- Unintentional/intentional poisoning
- Decline in food availability
- Electrocution and collision
- Habitat loss and degradation
- Wild capture and trade
- Persecution and hunting

CONSERVATION TIPS

- Do not disturb, approach or touch wildlife
- Talk about vulture conservation
- Do not fly drones close to nests or birds
- Support scientific research and monitoring
- Follow national laws and regulations

Environment Society of Oman

Funded by:
Disney Conservation Fund

* International Union for Conservation of Nature

© 2023

النسور المقيمة في عُمان

الرخمة

رخمة
Neophron percnopterus

التقديرات العالمية للأعداد: ١٨،٦٠٠ - ٥٤،٠٠٠ طائر، تتناقص
عدد الفصائل: ٣
تصنيف الاتحاد الدولي لصون الطبيعة: مهدد بالانقراض
الوضع في عُمان: مقيمين للتكاثر، بعض الزائرين في الشتاء

بيئة التكاثر

- التزاوج الأحادي
- التزاوج خلال الفترة من فبراير إلى مايو
- التعشيش على الجروف في الصدوع المظللة أو على الحواف والكهوف الصغيرة
- بقاء الزوجين معاً عبر السنين
- يضع الزوجين بيضتين اللتين تعطيان غالباً فرخين
- معدل الحضانة ٤٢ يوماً
- تبدأ الأفراخ بالطيران بعد ٧٠-٨٥ يوماً

سلوكيات التغذية

- تتغذى على الجيف ومخلفات الطعام في المرادم والمسالخ، وعلى بيوض الطيور أو براز الحيوانات أو صيد الطرائد الصغيرة
- يقوم الأبوين بإطعام الأفراخ عن طريق الارتجاع المعوي
- تتعلم الأفراخ من أبويها كيفية إيجاد الطعام ومهاراتها الاجتماعية

الحركة

- نطاق واسع
- هجرة جزئية
- تتكاثر في أوراسيا وأفريقيا، تُشتي الطيور المهاجرة في أفريقيا وجنوب آسيا

السلوكيات الاجتماعية

- محب للعشرة (اجتماعي)
- مناطق محددة للتزاوج
- تحطّ في مجموعات
- تحوم ضمن جماعات (بحثاً عن الغذاء)
- تتطفل على البشر (تستفيد من غذاء الناس)

النسر الآذن

النسر الآذن
Torgos tracheliotos

التقديرات العالمية للأعداد: ٩،٢٠٠ طائر، تتناقص
عدد الفصائل: ٢
تصنيف الاتحاد الدولي لصون الطبيعة: مهدد بالانقراض
الوضع في عُمان: مقيمين للتكاثر



بيئة التكاثر

- التزاوج الأحادي
- التزاوج خلال الفترة من نوفمبر إلى مايو
- التعشيش على قمم الأشجار المسطحة
- يتناوب الوالدان على حضانة الأفراخ ورعايتها وتظليلها
- يضع الزوجين عادةً بيضة واحدة
- معدل الحضانة ٥٥-٥٦ يوماً
- تبدأ الأفراخ بالطيران بعد ١٠٥ - ١٢٠ يوماً، وتبقى تحت إشراف الوالدين لمدة تزيد عن ٣ أشهر

سلوكيات التغذية

- تتغذى غالباً على الجيف، ولكنها قد تأكل الجراد والنمل الأبيض
- تزاحم بقية الأنواع وتغلبها على الجيف
- قادرة على هضم الجلد والأوتار والأربطة
- من المُمكن أن تفترس البيوض والأفراخ
- ترتاد على البقع المائية بشكل متكرر للشرب والاستحمام

الحركة

- نطاق واسع
- معظم الطيور البالغة غير مهاجرة باستثناء بعض الطيور خلال المواسم

السلوكيات الاجتماعية

- انعزالي إلى حدٍ كبير
- مناطق محددة للتزاوج
- تحطّ في مجموعات
- تحوم ضمن جماعات (بحثاً عن الغذاء)
- يطير الزوجين معاً غالباً بحثاً عن الطعام، وتحط (مرتفعات) معاً



هل تعلمون أن تصنيف نسور العالم الجديد يختلف عن نسور العالم القديم؟

التهديدات الرئيسية في عُمان

- التسمم المقصود/العرضي
- قلة موارد الطعام
- الصعق الكهربائي والتصادم
- تدهور الموائل الطبيعية وانحسارها
- الأسر والاتجار
- الصيد والقتل

نصائح للصون

- عدم إزعاج أو لمس أو مواجهة مفردات الحياة الفطرية
- انشروا الوعي حول صون النسور
- لا ترسلوا الكاميرات الطائرة قريباً من الطيور أو أعشاشها
- ادعموا الأبحاث والمسوحات العلمية
- التزموا بالقوانين والتشريعات المحلية

اكتشف المزيد

تعلموا المزيد عن الجوارح في عُمان وأوضاعها الراهنة هنا!

جمعية البيئة العمانية
Environment Society of Oman
بتمويلٍ من
Disney Conservation Fund

© ٢٠٢٣